

CAUTION

KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent



Tralkoxydim 400 WG

HERBICIDE

ACTIVE CONSTITUENT: 400 g/kg TRALKOXYDIM

GROUP A HERBICIDE

For the Control of Wild Oats and Annual Ryegrass and Suppression of Annual Phalaris in Wheat, Barley, Rye and Triticale.

IMPORTANT: Read this leaflet before use.

APVMA Approval No: 65110/54629

APPARENT PTY LTD A.C.N. 143 724 136

Suite G.08 | 762 Toorak Road, Hawthorn East VIC 3123 Australia
 T +61 3 9822 1321 F +61 3 9817 7845 E enquiries@apparentag.com.au
 www.apparentag.com.au

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply to weeds or crop under stress due to severe frost, drought, water logging, low soil fertility, pre-emergent herbicides, insects or diseases.

DO NOT use boomless jets or misters.

DO NOT apply if rain is expected within a half-hour.

Crop	Weeds Controlled	Weed Stage	State	Rate/ha	Critical Comments
Wheat, Barley, Rye, Triticale	Wild Oats	Growing actively at 2-4 leaf (pre-tillering)	Qld, NSW, Vic, Tas, SA only	300 g	Apply in first 4-6 weeks to unstressed crops and weeds when the majority of crop plants have 2 leaves. Application to lush crops may increase likelihood of yellowing and/or growth retardation.
				* 380 g	Apply in the first 4-6 weeks to unstressed crops and weeds when the majority of crop plants have 3 leaves. This is the minimum rate if tank mixing with listed broadleaf herbicides.
				* 430 g	Apply in first 4-6 weeks when the majority of crop plants have 3 leaves. Use this rate under dry soil conditions or if applying during August onwards. Use Magnify or Supercharge at 1 L per 100 L may improve weed control under these conditions.
	Wild oats and Annual Ryegrass in mixed infestations	Growing actively up to early tillering (Maximum of 2 tillers)	NSW, Vic, Tas, SA only	500 g	Use where application is delayed. Do not apply to pre-tillered crops. Use of Magnify or Supercharge at 1 L per 100 L may improve weed control if soil is dry or if applying during August onwards.
				* 380 g	Use when wild oats predominate. Do not apply to crops with less than 3 leaves. Use of Magnify or Supercharge at 1 L per 100 L may improve weed control if soil is dry or if applying during August onwards.
				* 500 g	Do not apply to pre-tillered crops. Use Magnify or Supercharge at 1 L per 100 L may improve weed control if soil is dry or if applying during August onwards.
Annual Phalaris (suppression only of Paradoxa grass, Lesser canary grass)	Growing actively at 2-4 leaf (pre-tillering)	NSW, Vic, Tas, SA only	380-500 g	Do not apply to tillered ryegrass. Use spray volumes to 70-100L/ha may assist control. Tank mixes of the lower rate with listed broadleaved herbicides may reduce ryegrass control.	
			* 380-500 g	Preferably apply within 6 weeks of sowing when weeds are at the 2 to 4 leaf stage and growing actively. Do not apply to tillered annual phalaris. The higher rate may give more reliable results. Use of spray volumes of 70-100 L/ha may assist control. Tank mixes with listed broadleaved herbicides may reduce annual phalaris control. Paradoxa grass is more sensitive than lesser canary grass. Do not apply to crops with less than 3 leaves.	
			* 380-500 g	Do not apply to tillered ryegrass. Use spray volumes to 70-100L/ha may assist control. Tank mixes of the lower rate with listed broadleaved herbicides may reduce ryegrass control.	
Wheat, Barley (except Moondyne Yagan), Rye, Triticale	Wild Oats	Growing actively at 2-4 leaf (pre-tillering)	WA only	250 g	Apply in first 4-6 weeks to unstressed crops and weeds when the majority of crop plants are at growth stage Z12. Application to lush crops may increase likelihood of yellowing and/or growth retardation.
				* 380 g	Use when application is delayed and the majority of crop plants are at growth stage Z13. This is the preferred rate for dense wild oat populations. Use this rate if tank mixing with listed broadleaf herbicides.
	Wild Oats and Annual Ryegrass in mixed infestations	Growing actively up to early tillering (Maximum of 2 tillers)	WA only	380 g	Use when wild oats predominate. Apply when the majority of crop plants are at growth stage Z13.
				* 380 g	Do not apply to tillered ryegrass. Use of higher spray volumes may assist control. Tank mixes with listed broadleaved herbicides may reduce ryegrass control.

*** ALWAYS APPLY MAGNIFY OR SUPERCHARGE. THE NORMAL RATE IS 750 mL PER 100 L SPRAY SOLUTION.**

INCREASE THE RATE TO 1 L PER 100 L UNDER DRYING CONDITIONS. ALWAYS USE 1 L PER 100 L FOR AERIAL APPLICATION.

NOT TO BE USED FOR ANY OTHER PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Resistant Weeds Warning

GROUP A HERBICIDE

Apparent Tralkoxydim 400 WG Herbicide is a member of the cyclohexanediones group of herbicides. Apparent Tralkoxydim 400 WG Herbicide has the inhibitors of acetyl CoA carboxylase mode of action. For weed resistance management Apparent Tralkoxydim 400 WG Herbicide is a Group A Herbicide. Some naturally occurring weed biotypes resistant to Apparent Tralkoxydim 400 WG Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Tralkoxydim 400 WG Herbicide or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Tralkoxydim 400 WG Herbicide to control resistant weeds.

MIXING

Only use sprayers in good working condition with high capacity agitation. Ensure the sprayer is cleaned to the label instructions on the product used prior to Apparent Tralkoxydim 400 WG Herbicide. Use only clean water for the spray solution. Ensure that all in-line strainer and nozzle screens in the sprayer are 100 mesh or coarser. DO NOT use cartridge or disc filters after the spray tank outlet, or have two (or more) filters directly after each other anywhere in the system.

Begin to fill spray with clean water, and engage agitator. Agitation must be continued throughout the entire mixing and spraying procedure.

When the sprayer is half full of water, add Apparent Tralkoxydim 400 WG Herbicide slowly to the spray tank through the tank opening or through a hopper designed to handle granules. The hopper should be dry before adding Apparent Tralkoxydim 400 WG Herbicide granules.

- Do not premix Apparent Tralkoxydim 400 WG Herbicide
- Add the total required amount of Apparent Tralkoxydim 400 WG Herbicide from all cartons first, prior to adding other herbicides or Magnify or Supercharge.
- Wait at least one minute after the last of Apparent Tralkoxydim 400 WG Herbicide has been added to the tank to allow for complete dispersion of the granules.
- Add the recommended broadleaf weed herbicide or insecticide next if tank mixing.
- Slowly add Magnify or Supercharge to the tank last, in at least 10 times its volume of water especially where a suction (filling) probe or hopper is used.
- Continue to fill tank to desired level with water.

SPRAYING INSTRUCTIONS

GROUND EQUIPMENT

Apply in a spray volume of 50-150 litres of water per hectare at a pressure of 250-350 kPa to ensure proper dispersion and spraying characteristics of the Apparent Tralkoxydim 400 WG Herbicide spray mix.

Use flat fan nozzles such as Spraying Systems 110-01, 110-015, 110-02 or equivalent sizes from other manufacturers. Do not use 110-03 or bigger nozzles with Apparent Tralkoxydim 400 WG Herbicide. Do not use floodjet nozzles, boomless jets or misters or controlled droplet application equipment. Nozzles maybe angled forward 45° to enhance penetration of the crop and provide better coverage. Always ensure that agitation is continued until spraying is completed even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes re-suspend the spray solution by running on full agitation until the settled particles are fully re-suspended, prior to spraying. The spray mixtures of Apparent Tralkoxydim 400 WG Herbicide and Magnify (or Supercharge) may be left overnight without loss of efficacy provided the settled product is completely re-suspended by stirring and/or agitation prior to re-use.

AERIAL EQUIPMENT

Apply through boom or Micronair units in 20-30 litres of water per hectare. The use of the higher volume is preferred due to more reliable results. Droplet size should be 200-250 microns (VMD) and swath width should not be wider than recommended for type of plane and application unit. Weed control following aerial application may not be as good as from ground application.

CROP SAFETY

If Apparent Tralkoxydim 400 WG Herbicide is applied to stressed crops or treatment is followed by stress or if crop plants have less than 2 true leaves (Z12), yellowing or growth retardation may occur. Crops normally recover without loss in yield. Avoid double spraying (overlap).

PLANTBACK PERIODS

Do not plant sorghum, maize, or rice for a period of four weeks after application of Apparent Tralkoxydim 400 WG Herbicide. Allow at least 10 days between application of Apparent Tralkoxydim 400 WG Herbicide and any other herbicide, except Ally for which the interval should be 14 days after Apparent Tralkoxydim 400 WG Herbicide. Allow at least 10 days between application of Apparent Tralkoxydim 400 WG Herbicide and any fertilizer treatment.

COMPATIBILITY

For broadleaved weed control, Apparent Tralkoxydim 400 WG Herbicide may be mixed as below.

State	Apparent Tralkoxydim g/ha	Co-herbicide maximum	Rate per ha
Qld, NSW, Vic, SA only	380 – 500 g	Bromoxynil + MCPA (200 + 200 g/L)	1.0 – 1.4 L
	380 – 500 g	MCPA ester	0.5/1.0 L
	380 – 500 g	Diffenofenican plus MCPA (25 + 250) (mixture may cause temporary chlorosis)	0.8 L
	380 – 500 g	Clopyralid (300 g/L)	0.3
WA only	380 g	Metosulam (714 g/kg)	7 g
	380 g	Bromoxynil + MCPA (200 + 200 g/L)	1.0 L
	380 g	MCPA ester * Note restriction on Kulin.	0.5 L *
	380 g	Diffenofenican plus MCPA (25 + 250) (mixture may cause temporary chlorosis)	0.8 L
	380 g	Clopyralid (300 g/L)	0.3 L
	380 g	Metosulam (714 g/kg)	7 g

DO NOT mix Apparent Tralkoxydim 400 WG Herbicide with 2,4-D, MCPA amine, dicamba or metsulfuron-methyl.

Allow at least 10 days before application of Apparent Tralkoxydim 400 WG Herbicide and herbicides containing 2,4-D, MCPA amine, dicamba or similar actives.

Allow at least 14 days between application of Apparent Tralkoxydim 400 WG Herbicide and metsulfuron-methyl.

Apparent Tralkoxydim 400 WG Herbicide should be applied prior to the broadleaved herbicides.

INSECTICIDES

Apparent Tralkoxydim 400 WG Herbicide is compatible with the emulsifiable formulations of the insecticides: permethrin, cypermethrin, phosmet, lambda-cyhalothrin and dimethoate.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Moderately hazardous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed original container in a dry cool well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilizers. Do not store for prolonged periods in direct sunlight.

Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty bags in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. If the product in eyes, wash it out immediately with water. When preparing the spray wear face shield or goggles. Wash hands after use. After each day's use wash face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which can be obtained from the supplier.

CONDITIONS OF SALE

Apparent Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Apparent Pty Ltd's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Apparent Pty Ltd has any authority to add to or alter these conditions.