

**DIRECTIONS FOR USE:****Restraints:**

**DO NOT USE on ewes which are producing or may in the future produce milk or milk products in human consumption or processing.**

**DO NOT USE on sheep with less than 6 weeks wool.**

**DO NOT collect and re-use fluid which runs off.**

**Contraindications: This product is contraindicated for use in animals with open wounds (e.g. those resulting from marking or mulesing operations).**

| ANIMAL/PEST                                         | TREATMENT                                               | INITIAL CHARGE | TOPPING UP     | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------------------|---------------------------------------------------------|----------------|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sheep Blowfly Control<br>( <i>Lucilia cuprina</i> ) | Jetting                                                 | 2 L per 1000 L | -              | <ol style="list-style-type: none"> <li>1. Jet at onset of blowfly wave.</li> <li>2. This product is not recommended for use in jetting races because a reduced period of protection may result unless the jetting race has been modified in accordance with agriculture NSW recommendations.</li> <li>3. Apply 2 to 4 L of jetting fluid per sheep, depending on sheep size and wool length.</li> <li>4. Ensure saturation to the skin.</li> <li>5. Pay particular attention to treatment of the crutch, pizzles and polls.</li> </ol>              |
|                                                     | Dipping<br>(Plunge – Shower and constant replenishment) | 2 L per 1000 L | 2 L per 1000 L | <ol style="list-style-type: none"> <li>1. Dip sheep on onset of fly wave.</li> <li>2. Thoroughly saturate sheep. When using a shower dip, run top and bottom sprays independently so as to ensure maximum pressure and therefore penetration.</li> <li>3. The product should not be used alone as a dip for blowfly control on sheep with less than 6 weeks wool as a reduced period of protection may result.</li> <li>4. The product is a non stripping dip. Reinforcement with undiluted concentrate is not required when topping up.</li> </ol> |
|                                                     | Dressing                                                | 2 L per 1000 L | -              | <ol style="list-style-type: none"> <li>1. Clip the wool surrounding the strike, treat the struck area and the surrounding wool with the recommended Diazinon knock down product (e.g. Top Clip® Blue Shield) then thoroughly jet the sheep with Independents Own Uranus Sheep Blowfly Treatment.</li> <li>2. This product is a slow killing larvicide when used on its own, larvae may take 3 to 4 days to die.</li> </ol>                                                                                                                          |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS**

Dipping sheep heavily infested with grass seeds is not recommended. Heavy rain following the application of this product could diminish the period of protection. If used in jetting races a reduced period of protection may result.

**MIXING****SHAKE WELL BEFORE USE.**

Settling may occur after storage, for some weeks. Stir product or invert container several times before opening. Pour the liquid into the vat through a strainer to remove any dry particles or flakes which can occasionally occur under hot storage conditions. Stir well. This product is readily soluble in both soft and hard water.

**COMPATIBILITY**

This product is compatible with Topdip® Blue Shield Sheep Dip, Di-Jet™, Topclip® Purple Shield Sheep Dip, 4 in 1 Sheep Dip®.

**MAINTENANCE OF DIPS**

1. Clean out dip, sump and spray lines thoroughly before charging. Disinfect with a registered bacteriostat at the manufacturer's recommended rate.
2. DO NOT use fouled wash. Shower dips should be emptied and cleaned out at the end of each day's dipping. Plunge dips should be emptied and cleaned out within 5 days of charging and more frequently if conditions favour post-dipping lameness infection, except in WA where plunge dips and sumps should be emptied daily.
3. Before use, stir well.

**JETTING**

Jet the sheep at onset of fly wave with efficient equipment which is able to

maintain the pressure of about 550 kPa for lambs and ewes in lamb.

**BACTERIOSTAT**

This product does not contain a bacteriostat. For control of post dipping infections a registered bacteriostat should be added at the manufacturer's recommended rate.

**WITHHOLDING PERIODS**

**MEAT: DO NOT USE less than 7 days before slaughter for human consumption.**

**MILK: DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption or processing.**

**WOOL HARVEST INTERVAL: DO NOT use less than 2 months before shearing or fibre collection.**

**TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):**

DO NOT use less than 21 days before slaughter for export. The ESI was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 03 9822 1321 or the APVMA website ([www.apvma.gov.au/residues/](http://www.apvma.gov.au/residues/)).

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

**SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container and preparing the dip or jetting fluid, wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, face shield and water resistant footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**Sheep Re-Handling Interval:** After treatment wait until sheep are dry before rehandling.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, rivers or streams with the chemical, discarded sump solution or used containers.

**DISPOSAL**

Triple or (preferably) pressure rinse into the dip vat. DO NOT dispose of undiluted chemicals on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Fouled or unwanted

sump solution should be placed in soil pit, away from dams, ponds and waterways. This container can be recycled if it is clean, dry, free of visible residues and has the drum/MUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinseate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drum/MUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

**STORAGE**

**KEEP OUT OF REACH OF CHILDREN.** Store below 30°C (Room Temperature), in original containers, tightly closed. DO NOT store for prolonged periods in sunlight.

**CONDITIONS OF SALE**

The use of Independents Own Uranus Sheep Blowfly Treatment being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

\* Other registered trademark

