

blank page

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Apparent** 

**Flutriafol 500 SC**

**FUNGICIDE**

ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL

**GROUP 3 FUNGICIDE**

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use.  
**IMPORTANT: Read this booklet before use.**

APVMA Approval No. 66731/54263

**APPARENT PTY LTD A.C.N. 143 724 136**  
Suite G.08 | 762 Toorak Road, Glen Iris VIC 3146 Australia  
M 0411 227 338 | E [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) [www.apparentag.com.au](http://www.apparentag.com.au)

054

**DIRECTIONS FOR USE: ALL STATES**

Crop	Disease	Rate per hectare	WHP
Canola	Blackleg ( <i>Leptosphaeria maculans</i> )	200 mL	4 weeks Grazing
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	100 or 200 mL	
	Septoria tritici blotch ( <i>S. tritici</i> ) Take-all ( <i>Gaeumannomyces graminis</i> )	200 mL	
Barley	Powdery mildew ( <i>Erysiphe graminis</i> ) Scald ( <i>Rhynchosporium secalis</i> )	100 or 200 mL	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS**

**APPLIED IN-FURROW AS A FERTILISER TREATMENT**

**CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING**

Critical Comments		
Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.		
<b>Application:</b> Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 200 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.		
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	
60	333*	
70	285	
80	250	
100	200	
120	166	
For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.		
<b>Application:</b> Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.		
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	
	100 mL rate	200 mL rate
50	200	400 *
60	166	333 *
70	143	285
80	125 †	250
100	100 †	200
120	88 †	166
* Application volumes over 300 mL per 100 kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.		
† Where application volume is low, product may be diluted with water to improve coverage.		

## GENERAL INSTRUCTIONS

### Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

**Sowing:** Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.

### FUNGICIDE RESISTANCE WARNING

#### GROUP 3 FUNGICIDE

Apparent Flutriafol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Apparent Flutriafol 500 SC Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Apparent Flutriafol 500 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Apparent Flutriafol 500 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Flutriafol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Apparent Pty Ltd should be advised immediately.

### SAFETY DIRECTIONS

Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length chemical resistant gloves and face shield or goggles. If product in eyes,

wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### SAFETY DATA SHEET

For further information refer to the safety data Sheet (SDS), which is available from the supplier.

### CONDITIONS OF SALE

The use of Apparent Flutriafol 500 SC Fungicide being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.