

DIRECTIONS FOR USE

Situation	Weeds Controlled	State	Rate L/ha	Critical Comments
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales	All States	4.5 in 330 – 550 L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
Transplanted	Feather, Shepherd's Purse, Nettles	Vic only	2.3 in 300 – 550 L	Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Apply on light sandy soils, low in organic matter.
Turf & Lawns Common Couch	Winter grass	All States	1.2 (12 mL/100m ²)	Rate/ Bowling Green 37 x 37m 170 mL
Queensland blue couch		Qld only	0.6 (6 mL/100m ²)	85 mL
Buffalo grass		WA only	1.2 (12 mL/100m ²)	170 mL
Lucerne, Clover, Medics etc. Growth for forage, Hay or Seed production	Annual Ryegrass, Winter grass, Barley grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue Wild Oats, Perennial Ryegrass, Bentgrass, English Couch, Yorkshire Fog Grass, Chickweed, seedling Sorrel, Blackberry Nightshade, Patterson's Curse, seedling Wireweed	NSW, ACT, SA, WA, Tas only	1 - 1.5 1.5 - 2	SEEDLING CROPS: Apply after the third trifoliolate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatments is recommended.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER

APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Apparent Piferer 500 Herbicide to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

RESISTANT WEEDS WARNING

GROUP D HERBICIDE

Apparent Piferer 500 Herbicide is a member of the benzimidazole group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management Apparent Piferer 500 Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to Apparent Piferer 500 Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Piferer 500 Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from failure of Apparent Piferer 500 Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be used can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Commission or Apparent Pty Ltd representative.

APVMA Approval No: 68267/102232

Batch No:

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and

skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of Apparent Piferer 500 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.



20L 9 345724 002314