

blank page

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## Turbo-Charged MCPA

### HERBICIDE

ACTIVE CONSTITUENTS: 275g/L TERBUTRYN  
160g/L MCPA present as the  
potassium salt

**GROUP C | I HERBICIDE**

For the control of seedling broadleaf weeds in wheat, barley, some varieties of oats, pasture and sugar cane as specified in the Directions for Use table.

**IMPORTANT: Read the attached booklet before use.**

APVMA Approval No: 69330/62660

**AIRR APPARENT PTY LTD**  
15/16 Princes Street, Newport NSW 2106  
M 0411 227 338 | [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) | [www.apparentag.com.au](http://www.apparentag.com.au)

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT apply to crops undersown with legumes.

DO NOT spray if weeds are wilted by dry or cold weather.

DO NOT spray if rain is expected within 6 hours.

DO NOT spray cereals or pastures on days when the temperature is greater than 22°C.

DO NOT add a wetting agent or a spraying oil for use in cereals or pasture.

Crop	Weeds Controlled
Wheat, barley and certain oat varieties②	Australian Crassula
	Cape Weed
	Charlock
	Fumitory-Red
	Fumitory-White
	Indian Hedge Mustard
	Melilotus/Hexam scent
	Paterson's Curse
	Rough Poppy
	Rounded Chamomile
	Sheepweed/Corn Gromwell
	Turnip Weed
	Wards weed
	Wild Turnip
Yellow Burr Weed	
Fat-hen	
Scented Woolly Sage	
Ball Mustard ①	
Deadnettle	
Mallee Catchfly	
Pimpernel	
Speedwell-lvyleaf	
Three-cornered Jack/ Spiny Emex/ Doublegee	
Toadrush	
Wireweed	
Saffron Thistle	
Stemless Thistle	

*Crassula sieberana*  
*Arctotheca calendula*  
*Sinapis arvensis*  
*Fumaria densiflora*  
*Fumaria parviflora*  
*Sisymbrium orientale*  
*Melilotus indica*  
*Echium plantagineum*  
*Papaver hybridum*  
*Matricaria matricarioides*  
*Buglossoides arvensis*  
*Rapistrum rugosum*  
*Carrichtera annua*  
*Brassica tournefortii*  
*Amsinckia* spp.

*Chenopodium album*  
*Salvia lanigera*

*Neslia paniculata*  
*Lamium amplexicaule*  
*Silene apetala*  
*Anagallis arvensis*  
*Veronica hederifolia*  
*Emex australis*  
*Juncus bufonis*  
*Polygonum aviculare*

*Carthamus lanatus*  
*Onopordum acaulon*

State	Rate/ha	Critical Comments
NSW, Vic, SA & WA only	1 L	Spray actively growing weeds at the 2 to 6 leaf stage when cereals have 3 to 5 leaves on the main stem, i.e. Zadok scale code Z13 - 15. Ensure yellow burr weed plants are at the 2 to 6 leaf stage. If plants are older than this use the higher rate. DO NOT spray this weed after the 12 leaf stage.
SA & WA only		
NSW, Vic, SA & WA only	1.5 L	Spray actively growing weeds at the 2 to 6 leaf stage when cereals have 3 to 5 leaves on the main stem, i.e. Zadok scale code Z13- 15. For Three-cornered Jack apply at the 1 to 4 true leaf stage. Control of lvyleaf Speedwell may at times be variable.
SA & WA only		

Crop	Weeds Controlled	
<b>Pastures</b> Use on Sub. clover (min 1 trifoliolate leaves) based pasture with either/ or Ryegrass, Phalaris, Cocksfoot (grasses min. 2 leaves)	Paterson's Curse (cotyledon to 2 leaf)	<i>Echium plantagineum</i>
<b>Pastures</b> Use on mixtures of either/or Sub. Clover (min. 3 trifoliolate leaves), Balansa clover (established plants), White clover (min. 5 trifoliolate leaves) with either/ or Ryegrass, Phalaris, Cocksfoot (grasses min. 3 leaves)	Capeweed (seedlings up to 20 cm diam.) Long Storksbill (seedlings up to 8 cm diam.) Paterson's Curse (seedlings up to 20 cm diam.) Toadrush (1 - 5 leaves) Crassula (1 - 5 cm diam.)	<i>Arctotheca calendula</i> <i>Erodium botrys</i> <i>Echium plantagineum</i> <i>Juncus bufonius</i> <i>Crassula</i> spp
Sugar cane (plant and ratoon)	Bellvine Morning glory Red convolvulus Pink convolvulus Star of Bethlehem Blackberry nightshade Red pigweed Giant (black) pigweed Pink burr Square weed Thornapple Sowthistle Native jute Calopo Rattlepod Annual ground cherry <b>Suppression only</b> Bluetop <sup>Ⓞ</sup> Common sensitive plant	<i>Ipomoea plebeian</i> <i>Ipomoea purpurea</i> <i>Ipomoea hederifolia</i> <i>Ipomoea triloba</i> <i>Ipomoea quamoclit</i> <i>Solanum nigrum</i> <i>Portulaca oleracea</i> <i>Trianthema portulacastrum</i> <i>Urena lobata</i> <i>Spermacoce latifolia</i> <i>Datura stramonium</i> <i>Sonchus oleraceus</i> <i>Corchorus trilocularis</i> <i>Calapogonium mucunoides</i> <i>Crotalaria</i> spp <i>Physalis ixiocarpa</i>  <i>Ageratum houstonianum</i> <i>Mimosa pudica</i>

State	Rate/ha	Critical Comments
NSW, Vic, SA, and WA only	300 mL to 500 mL	Apply soon after the break to the season when the weeds are up to the 2 leaf stage. Apply by boom spray or aircraft. Use the highest rate when the majority of plants are at the 2 leaf stage. Some reduction in clover growth may occur for a period after application. Plants usually recover and grow normally. Pasture should be growing well with adequate fertiliser and mite/insect control to help reduce the establishment of later germinating seedlings. Larger weeds present at spraying may not be controlled. <b>DO NOT</b> use on medics. <b>DO NOT</b> apply in less than 50 L/ha water. <b>DO NOT</b> use on seed crops. <b>DO NOT</b> spray after mid-August as damage may occur.
NSW, Vic, SA & WA only	1 L to 1.5 L	Apply by boom spray or aircraft. Use the highest rate when weeds are greater than 7 cm in diameter. Where a selective grass herbicide is to be used, apply Apparent Turbo-Charged MCPA prior to the application of the selective grass herbicide. Some reduction in clover growth may occur for a period after application. Plants usually recover and grow normally. <b>DO NOT</b> use on medics. <b>DO NOT</b> apply in less than 50 L/ha water. <b>DO NOT</b> use on seed crops. <b>DO NOT</b> spray after mid-August as damage may occur.
Qld, NSW, WA, NT only	2 L to 4 L <sup>Ⓞ</sup>	Apply post-emergent to broadleaf weeds up to 8 leaf or 30 cm diameter stage, vines to 1 m. Application should be made when weeds are growing actively and not suffering from moisture stress or waterlogging. Under cool conditions (<22° C) control may be slower and less effective. Can be applied over the top of cane. Some initial leaf yellowing and scorch may result but cane normally recovers well within 4 weeks. Caution should be exercised on new varieties of cane which may not have been tested. Use the lower rates on small weeds (less than 4 leaf) and the higher rates for larger weeds. Apply in a spray volume of 200 - 400 L/ha and ensure thorough coverage. Use the higher spray volumes where weed density is high or weeds are more advanced. <sup>Ⓞ</sup> Bluetop will only be suppressed. Use a minimum of 3 L/ha, on weeds less than 10 cm. Common sensitive plant - use a minimum of 3 L/ha

Crop	Weeds Controlled	
Sugar cane (plant and ratoon) (cont)	Green summer grass Crowsfoot grass Barnyard grass Guinea grass Summer grass Bluetop Centro Wild rose Bellvine Morning glory Red convolvulus Pink convolvulus Star of Bethlehem Giant sensitive plant Common sensitive plant Stinking passion vine Square weed Cinderella weed Giant (Black) pigweed Pink burr Thornapple Sowthistle Native jute Calopo Rattlepod Annual ground cherry Blackberry nightshade	<i>Brachiaria miliiformis</i> <i>Eleusine indica</i> <i>Echinochloa colona</i> <i>Panicum maximum</i> var <i>Hamil</i> <i>Digitaria ciliaris</i> <i>Ageratum houstonianum</i> <i>Centrosema pubescens</i> <i>Cleome aculeata</i> <i>Ipomoea plebeia</i> <i>Ipomoea purpurea</i> <i>Ipomoea hederifolia</i> <i>Ipomoea triloba</i> <i>Ipomoea quamoclit</i> <i>Mimosa invisa</i> <i>Mimosa pudica</i> <i>Passiflora foetida</i> <i>Spermacoce latifolia</i> <i>Synedrella nodiflora</i> <i>Trianthema portulacastrum</i> <i>Urena lobata</i> <i>Datura stramonium</i> <i>Sonchus oleraceus</i> <i>Corchorus trilocularis</i> <i>Calapogonium mucunoides</i> <i>Crotalaria spp</i> <i>Physalis ixiocarpa</i> <i>Solanum nigrum</i>

State	Rate/ha	Critical Comments
Qld, NSW, WA, NT only	3 L (+ 3 L ametryn 500 g/L) to 4 L (+ 3.7 L ametryn 500 g/L) ③	<b>Use in a tank mix with ametryn (500 g/L)</b> Apply between the early (1 - 4 leaf) and late (20 cm for grasses [Guinea grass and summer grass max. 15 cm] and 30 cm for broadleaf weeds) post-emergent stages of weeds. Use the lower rates on small weeds (less than 4 leaf) and for short residual control and the higher rates for large weeds and medium residual control. Where residual control is required the tank mix should be applied to moist soil and when rain or irrigation is expected within 10 days. Application should be made when weeds are growing actively and not suffering from moisture stress or waterlogging. Under cool conditions (<22°C) control may be slower and less effective. Apply in a spray volume of 200 - 400 L/ha and ensure thorough coverage. Use the higher volumes where weed density is high or weeds are more advanced. DO NOT apply this mixture by air. DO NOT apply over the top of cane. Use a directed spray to minimise contact with sugar cane foliage and shading of weeds. Continued from above <b>CAUTION:</b> Contact with the foliage of ametryn sensitive cane varieties such as Q119, Q122 and H56-752 should be avoided as they are susceptible to phytotoxicity from the tank mix. Caution should be exercised on new varieties of cane, which may not have been tested.

① Suppression only at 1.5 L/ha.

② Tolerant oat varieties are listed under the Crop Safety sub heading in the General Instructions section.

③ Add BS1000 120 mL/100 L or Agral 200 mL/100 L (sugar cane only)

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**DO NOT GRAZE BARLEY, WHEAT, OATS OR PASTURE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**DO NOT APPLY THE TANK MIX OF APPARENT TURBO-CHARGED MCPA AND AMETRYN (500 g/L) LATER THAN 9 MONTHS BEFORE HARVEST.**

## GENERAL INSTRUCTIONS

### RESISTANT WEEDS WARNING

#### GROUP C I HERBICIDE

Apparent Turbo-Charged MCPA Herbicide is a member of both the triazines and phenoxy's groups of herbicides. Apparent Turbo-Charged MCPA has the inhibitor of photosynthesis at photosystem II and disruptor of plant cell growth modes of action. For weed resistance management Apparent Turbo-Charged MCPA is a group C and I herbicide.

Some naturally occurring weed biotypes resistant to Apparent Turbo-Charged MCPA Herbicide and other group C or I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Turbo-Charged MCPA Herbicide or other group C or I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Turbo-Charged MCPA Herbicide to control resistant weeds.

### MIXING

Apparent Turbo-Charged MCPA Herbicide mixes readily with water. Half fill the spray tank with water. Shake the container several times before use and add the recommended quantity to the tank. Keep the spray mixture well agitated during mixing and spraying using either mechanical or venturi type agitation that has sufficient water movement for the solution to remain in suspension. Reseal part-used containers immediately. Wash sprayer thoroughly with clean water after use.

### APPLICATION

**Boom Sprayer:** Calibrate the sprayer before use to ensure the correct application rate is applied per hectare. Check boom height for even application.

**Cereal and pastures:** Apply 50 to 100 L of water/ha using no smaller than 80 mesh pump and jet filters. Use the higher volume if the weed infestation is heavy or the crop cover is dense.

**Sugar cane:** Apply 200 to 400 L of water/ha using no smaller than 80 mesh pump and jet filters. Use the higher volume if the weed density is high or on more advanced weeds. Thorough coverage is necessary. Always add the required wetter.

**Aircraft:** Apply 20 to 30 L of water/ha. The higher volume is preferred due to more reliable results. Weed control following aerial application may not be as good as from ground application.

### COMPATIBILITY

This product can be tank mixed with metsulfuron methyl herbicide for the control of specific weeds such as capeweed, sheepweed and toadrush. DO NOT tank mix with diclofop methyl or tralkoxydim as significant crop injury and reduced weed control will occur. In pastures this product can be mixed with MCPA amine and the insecticides dimethoate, lambda-cyhalothrin, alpha-cypermethrin and omethoate (ensure good agitation is maintained with this product and avoid use of cold water). In sugar cane this product can be tank mixed with Apparent Wetter 600, Apparent Wetter 1000, 2,4-D amine and ametryn (500 g/L).

When tank mixing read both labels and adhere to the crop recommendations and directions that are most restrictive. The compatibilities given are based on experience and are considered correct. However, as changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, AIRR Apparent Pty Ltd accepts no responsibility for the behaviour of such mixtures.

### TIMING

Weeds should be sprayed while actively growing and at the 2 to 6 true leaf stage when they are most susceptible. As a general guide Apparent Turbo-Charged MCPA Herbicide should be applied approximately 5 to 8 weeks after planting. Extremes of cold or drought are unfavourable. The most favourable conditions are when soil is moist, weather fine and rain unlikely within 6 hours following application. In pasture, weeds should be up to 20 cm diameter but refer to the directions for use table for specific details. In sugar cane, apply Apparent Turbo-Charged MCPA Herbicide alone to broadleaf weeds up to the 8 leaf or 30 cm diameter stage, vines to 1 m length.

### CROP SAFETY

Research has shown that although some early growth check and leaf margin scorch can occur in oats, if grain yield is measured, the following varieties are tolerant or moderately tolerant to Apparent Turbo-Charged MCPA Herbicide.

**Tolerant:** (apply up to 1.5 L/ha) Bandicoot, Coolabah, Mortlock, Potaroo, Swan, Wallaroo, West.

**Moderately Tolerant:** (DO NOT exceed 1 L/ha) Dalyup, Windjardie.

DO NOT spray other oat varieties unless you seek advice before spraying.

In pastures sub. clover growth may be reduced for a period after application but plants usually recover and spring growth is not normally affected.

Sugar cane varieties such as Q122 may show yellowing and some leaf scorch immediately after spraying but plants normally recover within 4 weeks.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers, or waterways with the chemical or used containers.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated locked area away from children, animals, food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple or, preferably, pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this

purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SAFETY DIRECTIONS**

Harmful if swallowed. Attacks eyes. Will irritate the skin. Avoid contact with skin. Repeated exposure may cause allergic disorders. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia:13 11 26.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

### **CONDITIONS OF SALE**

The use of Apparent Turbo-Charged MCPA Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AIRR Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.