

Apparent Pty Ltd
Suite G.08, 762 Toorak Road,
Glen Iris VIC 3146,
PO Box 3092 Cotham PO
Kew Vic 3101
Phone 03 9822 1321 Mobile 0411 227 338

APVMA Approval No: 69913/62050

CAUTION

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Independents Own

Fenbender 100

Oral Anthelmintic for cattle
& horses

ACTIVE CONSTITUENT:
100 g/L FENBENDAZOLE

For the control of benzimidazole sensitive roundworms and lungworms in cattle, and large and small strongyles, ascarids, and pinworms in horses.

**IMPORTANT: READ THIS LEAFLET
BEFORE USING THIS PRODUCT**

MANUFACTURED BY WSD

Koojan Avenue, South Guildford WA 6055
ABN 58 009 372 525

PROUDLY



MADE &
OWNED

iO FENBENDER 100 ORAL ANTHELMINTIC FOR CATTLE AND HORSES:

Contains Fenbendazole, a member of the benzimidazole family of drenches. It is effective in cattle and horses against benzimidazole sensitive strains of the following internal parasites:

CATTLE

Barber's pole worm	<i>Haemonchus placei</i>
Small brown stomach worm	<i>Ostertagia</i> spp.
Small hair worm	<i>Trichostrongylus</i> spp.
Nodule worm	<i>Oesophagostomum radiatum</i>
Thin necked intestinal worm	<i>Nematodirus</i> spp.
Small intestinal worm	<i>Cooperia</i> spp.
Lungworm	<i>Dictyocaulus viviparus</i>
Whipworm	<i>Trichuris</i> spp.
	<i>Capillaria</i> spp.

HORSES

Large strongyles	<i>S. vulgaris</i> , <i>S. edentatus</i> , <i>S. equinus</i>
Small strongyles	<i>Trichonema</i> spp., <i>Triodontophorus</i> spp.
Ascarids	<i>Parascaris equorum</i>
	<i>Oxyuris</i> spp.

DIRECTIONS FOR USE

DOSAGE AND ADMINISTRATION SHAKE WELL BEFORE USE.

iO Fenbender 100 can be administered orally through standard accurate drenching equipment. Clean equipment thoroughly before and after use.

CATTLE: 3.75 mL/50 kg bodyweight (7.5 mg Fenbendazole per kg bodyweight).

Bodyweight CATTLE	Dose Rate Bodyweight	10 L Treats	5 L Treats	1 L Treats
50 kg	3.75 mL	2666	1333	266
100 kg	7.50 mL	1333	666	133
150 kg	11.25 mL	888	444	88
200 kg	15.00 mL	666	333	66
250 kg	18.75 mL	533	266	53
300 kg	22.50 mL	444	222	44
350 kg	26.25 mL	380	190	38
400 kg	30.00 mL	333	166	33
450 kg	33.75 mL	296	148	29
500 kg	37.50 mL	266	133	26
550 kg	41.25 mL	242	121	24
600 kg	45.00 mL	222	111	22
650 kg	48.75 mL	205	102	20
700 kg	52.5 mL	190	95	19
750 kg	56.25 mL	177	88	17
Over 750 kg	3.75 mL per 50 kg			

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

HORSES: 1 mL/10 kg bodyweight (10 mg Fenbendazole per kg bodyweight).

For the control of *Strongyloides westeri* in foals:

5 mL/10 kg bodyweight (50 mg Fenbendazole per kg bodyweight).

Bodyweight HORSES	Dose Rate Bodyweight	10 L Treats	5 L Treats	1 L Treats
100 kg	10 mL	1000	500	100
150 kg	15 mL	667	333	67
200 kg	20 mL	500	250	50
250 kg	25 mL	400	200	40
300 kg	30 mL	333	167	33
350 kg	35 mL	286	143	29
400 kg	40 mL	250	125	25
450 kg	45 mL	222	111	22
500 kg	50 mL	200	100	20
550 kg	55 mL	182	91	18
600 kg	60 mL	167	83	17
650 kg	65 mL	154	77	15
700 kg	70 mL	143	71	14
750 kg	75 mL	133	67	13
Over 750 kg	1 mL per 10 kg			

Internal parasites may develop resistance to drenches (anthelmintics).

Ask your local veterinary adviser for recommended parasite management practices for your area to reduce development of resistance.

It is advisable that a drench resistance test be conducted before any drench is used. If drench failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer.

WITHHOLDING PERIODS

MEAT (Horses): DO NOT USE less than twenty eight (28) days before slaughter for human consumption.

MEAT (Cattle): DO NOT USE less than fourteen (14) days before slaughter for human consumption.

MILK: Zero (0) Days.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than fourteen (14) days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Apparent Pty Ltd Phone 03 9822 1321 or the APVMA website www.apvma.gov.au/residues/.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. Do not contaminate streams, rivers and waterways with this product or used containers.

DISPOSAL 1L pack size

Dispose of empty container by wrapping with paper and putting in garbage.

5L Backpack, 5L, 10L, pack sizes

Triple or (preferably) pressure rinse. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned.

STORAGE

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area, and protect from light.

CONDITIONS OF SALE

The use of Independents Own Fenbender 100 Oral Anthelmintic for Cattle and Horses being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used

by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.