

blank page

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Apparent** 

## Decimator 400

### INSECTICIDE

ACTIVE CONSTITUENT: 400 g/L DIMETHOATE  
(an anticholinesterase compound)

**GROUP 1B INSECTICIDE**

For the control of a wide range of insect pests on certain fruit trees and vegetables, citrus, pastures, cotton, lucerne, peanuts and ornamentals as listed in the Directions for Use Table.

**IMPORTANT: Read the attached booklet before use.**

APVMA Approval No: 70165/RV0317D

**APPARENT PTY LTD A.C.N. 143 724 136**  
Suite G.08 | 762 Toorak Road, Glen Iris VIC 3146 Australia  
M 0411 227 338 | E [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) [www.apparentag.com.au](http://www.apparentag.com.au)

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT use to control pests that are resistant to organophosphorus insecticides as treatment may be ineffective.

DO NOT apply to any non-food tree crop or plantation (including Eucalyptus spp.) by air.

DO NOT apply by misting or fogging equipment

DO NOT apply with air blast spray equipment unless operators are protected by engineering controls such as enclosed cabs fitted with appropriate air filters

DO NOT use open mixing/loading systems for aerial application

**FIELD CROPS**

CROP	PEST	STATE	RATE	WHP (days)
<b>Cereals</b> (Wheat, Barley, Oats, Triticale)	Lucerne Flea	NSW, Vic, Tas, SA, WA only	55 - 85 mL/ha	4 weeks (harvest)
	Red legged earth mite	Vic, Tas, SA, WA only		14 days (grazing)
		NSW only	85 mL/ha	
		NSW, Vic, Tas, SA, WA only	200 mL/ha	
	Wingless Grasshopper	All States	75 mL/100 L of water or 750 mL/ha	
	Brown Wheat Mite	Qld, WA only	90 mL/ha	
	Blue Oat Mite	Qld, NSW, WA only		
Leafhoppers, Cereal Aphids	All States	500 mL/ha		
<b>Pastures</b> <b>Pasture Seed and Forage Crops, (Inc. Clover, Medics, Cereals, Lucerne, Legumes for animal feed)</b>	Lucerne Flea, Redlegged Earth Mite	NSW, Vic, Tas, SA, WA only	55 - 85 mL/ha	-
	Redlegged Earth Mite		250 mL/ha	
<b>Lucerne</b>	Lucerne Flea	NSW, Tas, Vic, SA, WA only	55 - 85 mL/ha	
	Redlegged Earth Mite	SA, Tas, Vic, WA only	55 - 85 mL/ha	
		NSW only	85 mL/ha	
<b>Maize</b>	Maize Leafhoppers, Thrips	Qld, WA only	500 mL/ha	4 weeks harvest 14 days grazing
<b>Sorghum</b>	Aphids	Qld, WA only	500 mL/ha	4 weeks (H) 14 days(G)
<b>Tobacco</b>	Lucerne Flea, Redlegged Earth Mite	NSW, WA only	80 mL/100 L of water	4 weeks
<b>Adzuki Beans, Cowpeas, Mung Beans, Navy Beans, Pigeon Peas, Chickpeas, Lupins, Borlotti Beans</b>	Aphids (excluding Green Peach Aphid)	All States	500 mL/ha	14 harvest 14 grazing
	Mirid Bugs			
	Thrips (including Bean Blossom Thrips) (except in Qld cowpeas), Bean Fly, Leafhoppers (including Jassids), Green Peach Aphid		800 mL/ha or 75 mL/100 L of water	
<b>Field Peas and Beans</b>	Aphids, Thrips, Leafhoppers (including Jassids), Mites (including Spider mites), Bugs (including Green Vegetable Bug, Bean Fly, Red Legged Earth Mite		75 mL/100 L of water or 800 mL/ha	

CRITICAL COMMENTS
DO NOT harvest for 4 weeks after application. DO NOT graze or cut for stock feed for 14 days after application. Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. DO NOT spray on bare ground. Allow the crop to emerge before application. Apply from boom spray in 50 - 100 L water/ha or Aircraft and misting machines in 20 - 40 L of water per hectare.
A well timed application at this rate may provide an extended period of control. Apply as above. See General Instructions.
Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
Apply when pests appear.
Apply when pests threaten to damage crop.
Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in NSW and in cold weather and/or for heavy infestations in other States. DO NOT spray on bare ground. Allow the crop to emerge before application. <b>Boom spray:</b> Apply in 50 - 100 L of water/ha. <b>Aircraft:</b> Apply in 20 - 40 L/ha
A well timed application at this rate may provide an extended period of control. Apply as above. See General Instructions.
Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather or on mature pastures. DO NOT spray on bare ground. Allow the lucerne to emerge before application. DO NOT use more than 7 days after crop emergence. <b>Boom spray:</b> Apply in 50 - 100 L of water/ha. <b>Aircraft:</b> Apply in 20 - 40 L/ha
DO NOT harvest for 4 weeks after application. DO NOT graze or cut for stock feed for 14 days after application Apply 2 sprays 5 - 7 days apart.
DO NOT harvest for 4 weeks after application. DO NOT graze or cut for stock feed for 14 days after application Apply as required.
Apply spray to tobacco in seedbed when insects are present. Re-apply after 7 days if necessary.
DO NOT harvest for 14 days after application. DO NOT graze or cut for stock feed for 14 days after application Apply when flower spikes carry 20 to 50 aphids and repeat as necessary Apply when insects appear and repeat as necessary. DO NOT re-apply within 14 days
For Thrips (excluding Bean Blossom Thrips): Two treatments between pre-bloom and pod initiation may be necessary. Apply both sprays early during this period if infestation is severe or prolonged. Use sufficient water to give good coverage. For Bean Fly, Bean Blossom Thrips and Leafhoppers: Apply when pests appear. For Green Peach Aphid: Apply when flower spikes carry 20 to 50 aphids and repeat as necessary. DO NOT re-apply within 14 days
Apply when pests appear and repeat as necessary. For Green vegetable bug apply in first flowering and repeat 3 weeks later. DO NOT re-apply within 14 days

CROP	PEST	STATE	RATE	WHP (days)
<b>Lentils</b>	Redlegged Earth Mite	All States	90 mL/ha	14 harvest
<b>Soy Beans</b>	Green Vegetable Bug, Leafhoppers (including Jassids)		340 mL/ha	
<b>Grain Legumes</b>	Spider Mites, Thrips, Jassids, Green Vegetable Bug, Aphids, Bean Fly	Qld, Vic, Tas, SA, WA only	75 mL/100 L or 800 mL/ha	14 grazing
	Redlegged Earth Mite	Vic, Tas, SA, WA only	75 mL/100 L	
	Lucerne Flea	WA only	85 mL/100 L	

#### OILSEED AND FIBRE CROPS

CROP	PEST	STATE	RATE	WHP (days)	
<b>Oil Seeds other than peanuts and cotton (including Mustard, Linseed, Peanut, Poppy, Canola, Safflower, Sunflower)</b>	Lucerne Flea	NSW, Vic, Tas, SA only	55 - 85 mL/ha	-	
		WA only	40 - 55 mL/ha		
	Redlegged Earth Mite	Vic, Tas, SA, only	55-85 mL/ha		
		WA only	40 - 55 mL/ha		
<b>Cotton</b>	Aphids, Spider Mites, inc. Red Spider Mite, Two Spotted Mite	NSW, Qld, WA only	500 mL/ha	Harvest 14	
			Thrips		350 to 375 mL/ha
	Wingless Grasshoppers	All States	750 mL/ha or 75 mL/100 L of water	Harvest 14	
	Leafhoppers (including Jassids), Green Vegetable Bug	All States	350 mL/ha		
	Bugs, inc. Green Mirids, Broken Backed Bug, Apple Dimpling Bug, Brown Smudge Bug, Rutherglen Bug	NSW, Qld, WA only	340 to 500 mL/ha		
	Lucerne Flea	NSW, Vic, Tas, SA only	55 - 85 mL/ha		
		WA only	45 - 55 mL/ha		
	Red legged earth mite	Vic, Tas, SA only	55 - 85 mL/ha		
		WA only	45-55 mL/ha		
		NSW, only	85 mL/ha		
<b>Peanuts</b>	Aphids, Jassids, Green Vegetable Bug	All States	350 mL/ha		Harvest 14
	Peanut Mite, Thrips	Qld, NSW, WA only	750 mL/ha or 75 mL/100 L of water		Grazing 14
	Wingless Grasshoppers	All States			
	Lucerne Flea	NSW, Vic, Tas, SA only	55 - 85 mL/ha	Harvest 14	
		WA only	45 - 55 mL/ha		
	Red legged earth mite	Vic, Tas, SA only	55 - 85 mL/ha		
WA only		45-55 mL/ha			
NSW, only		85 mL/ha			

#### FRUIT CROPS

CROP	PEST	STATE	RATE	WHP (days)
<b>Berry Fruits (Blackberries, Raspberries ONLY)</b>	Spider Mites, Thrips, Jassids, Aphids, Red legged Earth Mite	All States	75 mL/100 L of water	7
	Strawberry Bug, Rutherglen Bug	QLD, Vic, Tas, SA, WA only		

CRITICAL COMMENTS
Apply when pests appear. DO NOT re-apply within 14 days
DO NOT harvest for 14 days after application. DO NOT graze or cut for stock feed for 14 days after application. Apply when insects appear and repeat as necessary. Spray when flowering spikes carrying 20 - 50 aphids are easy to find and when there is evidence of viral disease. Some strains of Spider Mite are resistant to organophosphorus compounds.
Apply at emergence.

CRITICAL COMMENTS
Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. DO NOT spray on bare ground. Allow the crop to emerge before application. Boom spray: Apply in 50 - 100 L of water/ha. Aircraft: Apply in 20 - 40 L/ha.
DO NOT harvest for 14 days after application. DO NOT graze or cut for stock feed. DO NOT feed cotton fodder, stubble or trash to livestock. Apply when pests appear and repeat as required. Use the higher rate for heavy infestations. Some strains of Spider Mite are resistant to organophosphorus compounds. DO NOT use this product where resistant strains are present.
Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. DO NOT spray on bare ground. Allow the crop to emerge before application. Boom spray: Apply in 50 - 100 L water/ha. Aircraft: Apply in 20 - 40 Lha.
DO NOT harvest for 14 days after application. DO NOT graze or cut for stock feed for 14 days after application. Apply when pests appear.
Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
Apply 3 - 5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. DO NOT spray on bare ground. Allow the crop to emerge before application. <b>Boom spray:</b> Apply in 50 - 100 L water/ha. <b>Aircraft:</b> Apply in 20 - 40 Lha.

CRITICAL COMMENTS
Apply when pest first appears and repeat at 3 weekly intervals or as necessary. Some strains of Spider Mites are resistant to organophosphorus compounds.

CROP	PEST	STATE	RATE	WHP (days)
Blueberries, Bilberries, and other Vaccinium Berries	Queensland Fruit Fly	NSW, WA only	75 mL/100 L of water	1
	Spider mites, Thrips, Jassids Aphids, Red legged earth mite	All states		
	Strawberry Bug, Rutherglen Bug	QLD, Vic, Tas, SA, WA only		

#### FRUIT TREES

RATE				
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Application Section.				
CROP	PEST	STATE	RATE	WHP (days)
Avocados	Queensland Fruit Fly	Qld, WA, NT only	75 mL/100 L as an overall spray	7
Citrus Fruit (including Oranges, Lemons, Mandarins, Limes) (except Meyer Lemons, Seville Oranges and Cumquats)	Queensland Fruit Fly	Qld, NSW, Vic, WA only	75 mL/100 L of water	7
	Mediterranean Fruit Fly	WA, Vic only	75 mL/100 L of water	
	Aphids, Thrips	All States		
	Bronze Orange Bug	Qld, NSW, Vic, SA, WA only		
Wingless Grasshopper	All States			
Litchi	Litchi Erinose Mite	Qld, NSW, WA only	75 mL/100 L of water	-
Mangoes	Queensland Fruit Fly	Qld, NSW, Vic, WA, NT only	75 mL/100 L of water	7
	Mediterranean Fruit Fly	NSW, Vic, WA, only		3

#### VEGETABLES

<b>Vegetables: Use ONLY on the following:</b>	Aphids, Jassids, Mites, Leaf Hoppers, Green Vegetable Bug, Thrips, Wingless Grasshoppers	All States	75 mL/100 L of water (or 750 mL/ha for Wingless Grasshoppers)				
Tomatoes, large, field grown for fresh consumption				Not required when used as directed			
Melons, Zucchini				1 (Zucchini) 7 (Melons)			
Capsicums,				3			
Asparagus, Onions, Rhubarb, Sweetcorn				7			
Beans, Peas (Not snow or sugar snap peas)				7 (H, G)			
Beetroot, Eggplant, Potatoes, Sweet Potatoes, Turnip				14			
Tomatoes for processing				21			
Beans, Peas				Cow Pea Aphid	NSW, WA only	350 - 650 mL/ha	7 (H, G)
				Bean Fly	All States	75 mL/100 L of water or 750 mL/ha	
				Redlegged Earth Mite	NSW, Vic, Tas, SA, WA, only	800 mL/ha or 75 mL/100 L of water	
Beetroot	Leafmining Fly	NSW only	800 mL/ha or 75 mL/100 L of water	14			

CRITICAL COMMENTS
DO NOT exceed a maximum number of 7 applications per crop per season with a minimum retreatment interval of 21 days between consecutive applications. DO NOT harvest for 1 day after final application.

CRITICAL COMMENTS
For all tree and vine crops in this table: Apply by dilute or concentrate spraying equipment. For concentrate spraying, refer to the Application Section. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CRITICAL COMMENTS
Apply as pest populations indicate.
QLD, NSW, VIC ONLY: Do not use on Meyer Lemons, Seville Oranges and Cumquats. Apply two full cover sprays 2 weeks apart, 7 weeks and 5 weeks before harvest. If harvesting is delayed a third spray may be required. WA ONLY: Apply about 6 weeks before fruit ripens. Reapply at fortnightly intervals. The last spray should be one week before fruit ripens.
Apply when pests appear.
Apply when pest appears and repeat as necessary.
Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
Pre-planting Dip: Immerse plants in mixture for 1 minute and drain before planting in the field.
Established trees: Apply just before a growth flush and repeat at 14 - 21 day intervals until all new growth is damage free.
Apply as a cover spray at first sign of infestation.

Apply when pests appear. This product will not control OP resistant mites.
<b>Tomatoes, large, field grown for fresh consumption:</b> DO NOT apply after commencement of flowering; DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, green houses or plastic tunnels; DO NOT USE on cherry, grape or mini tomatoes.
<b>Wingless Grasshoppers:</b> Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
<b>Tomatoes for processing:</b> DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, green houses or plastic tunnels; DO NOT USE on cherry, grape or mini tomatoes.
Apply when pests appear. Use the higher rate in cold weather.
Apply when pest damage first appears. Repeat spray if necessary.

**VEGETABLES (cont)**

CROP	PEST	STATE	RATE	WHP (days)
<b>Capsicums</b>	Cucumber fly	NSW, WA only	75 mL/100 L of water or 750 mL/ha	3
	Fruit fly			
<b>Cucurbits: Zucchini and melons ONLY.</b>	Cucumber Fly	Qld, NSW, WA, NT only		1 (Zucchini) 7 (Melons)
<b>Tomatoes (for processing ONLY)</b>	Queensland Fruit Fly	Qld, NSW, Vic, WA, only	75 mL/100 L of water or 750 mL/ha	21
	Mediterranean Fruit Fly	NSW, Vic, WA only		
	Tomato Mite	NSW, Vic, Tas, SA only	60 mL/100 L	
	Bryobia Mite	Vic, Tas, SA, WA only		
<b>Tomatoes, large, field grown for fresh consumption</b>	Tomato Mite	NSW, Vic, Tas, SA only	60 mL/100 L	Not required when used as directed
	Bryobia Mite	Vic, Tas, SA, WA only		
<b>Beetroot, Onions</b>	Redlegged Earth Mite	NSW, Vic, Tas, SA, WA only	75 mL/100 L of water	14 7

**POST HARVEST DIPPING**

CROP	PEST	STATE	RATE	WHP (days)
<b>Avocados, Chinese Gooseberries (Kiwifruit) (inedible peel varieties ONLY), Lychees,</b>	Queensland Fruit Fly	NSW, WA only	Charge the dip at a rate of 100 mL/100 L of water	-
<b>Bananas</b>	Fruit Fly	NSW, WA only	75 mL/100 L water	-
<b>Custard apple</b>	Queensland Fruit Fly	NSW, WA, NT only	Charge the dip at a rate of 100 mL/100 L of water	-
<b>Mangoes, Pawpaws, Passionfruit</b>		NSW, WA, only		-

**POST HARVEST DIPS – NOTE THIS IS A QUARANTINE TREATMENT ONLY**

CROP	PEST	STATE	RATE	WHP (days)
<b>Avocados, Bananas, Cactus Fruit, Custard Apples, Feijoa, Guavas (inedible peel varieties ONLY), Kiwifruit (inedible peel varieties ONLY), Mangoes, Pawpaws, Banana Passionfruit, Passionfruit, Pomegranate, Tamarillos</b>	Queensland Fruit Fly ( <i>Dacus tryoni</i> )	Qld, NSW, WA, NT only	100 mL/100 L of water	-
<b>Mangoes</b>	Darwin Fruit Fly ( <i>Bactrocera aquilonis</i> )	WA, NT only		

CRITICAL COMMENTS
Apply when insects appear. DO NOT USE as a post-harvest or post-harvest quarantine treatment
Apply when pests first appear and repeat as required.
Apply when pests appear and repeat as required.
QLD ONLY: Apply two full cover sprays 4 weeks before harvest. NSW ONLY: Apply two full cover sprays 4 weeks and 3 weeks before harvest. Vic only: Apply at 7 and 5 weeks before harvest. WA ONLY: Apply about 6 weeks before fruit ripens. The last spray should be three weeks before harvest. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, green houses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes.
Apply as a cover spray 4 weeks before harvest. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, green houses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes.
Apply as a cover spray. DO NOT apply after commencement of flowering. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, green houses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes.
Apply when pests first appear and repeat at 3 weekly intervals as required.

CRITICAL COMMENTS
Dip the fruit for 1 minute and allow to drain before packing.
Dip fruit for 10 - 60 seconds. Top with concentration of 125 mL - 150 mL/100 L.
Dip the fruit for 1 minute and allow fruit to drain before packing.

CRITICAL COMMENTS
DIPPING: Immerse produce in emulsion for 1 minute or according to the requirements of the importing State or Country. TOPPING UP: (400 ppm dimethoate emulsion only): Top up with a separately prepared 400 ppm (100 mL/100 L) emulsion. REINFORCEMENT: (400 ppm dimethoate emulsion only): After each week, add 3 mL of product/100 L of dip emulsion. NOTE: 1. Refer also to Refnote R6/Feb 83 (Agdex 201/681) – "FRUIT AND VEGETABLES- stability of dimethoate in dips". 2. 400 ppm is the dip concentration required for fruit fly susceptible produce destined for interstate markets. For other destinations the requirements may differ (eg. fruit for export to New Zealand to be treated at 500 ppm); check with relevant authorities.

**MISCELLANEOUS**

CROP	PEST	STATE	RATE	WHP (days)	CRITICAL COMMENTS
<b>Ornamentals (not Chrysanthemum, Begonias, Liquid Amber or Gloxinias)</b>	Aphids, Thrips, Jassids, Spider Mites, Leafhoppers, Azalea Lace Bug, Green Vegetable Bug, Leaf Miners, Greenhouse White Fly, Wingless Grasshopper	All States	75 mL/ 100 L of water	-	Apply when pests appear and repeat as necessary. Some strains of Spider Mites are resistant to organophosphorus compounds. <b>Wingless Grasshoppers:</b> In addition to the infested area spray a band of about 20 metres around areas to be protected.
	Bronze Orange Bug	Qld, NSW, Vic, SA, WA only			
	Woolly Aphid	Vic, Tas, SA, WA, NT only			
<b>Ornamental Shrubs</b>	Sap-sucking and Leaf-eating insects (including Aphids, Mites, Leafhoppers (including Jassids), Mealybugs, Sawflies, Leafminers, White Flies, Wingless Grasshopper, Psyllids, Scales, Scarab and Leaf Beetles and Beetle Larvae, Moth Caterpillars, Lace Bugs, Gall Insects), Azalea Lace Bug, Green Vegetable Bug, Rutherglen Bug	All States	75 mL/100 L water	-	Apply when pests first appear ensuring thorough coverage of foliage. Repeat as required. Apply late afternoon to prevent burning of foliage and to avoid affecting foraging birds and beneficial insects. DO NOT spray prior to or during rain. Avoid spray drift. DO NOT harvest fruit or other produce from sprayed trees. <b>DO NOT use on Chrysanthemums, Begonias, Liquidamber or Gloxinias.</b>
	<b>Ornamental Farm and Forest Trees</b>	Wingless Grasshopper, Psyllids, Scales, Scarab and Leaf Beetles and Beetle Larvae, Moth Caterpillars, Lace Bugs, Gall Insects), Azalea Lace Bug, Green Vegetable Bug, Rutherglen Bug	WA only	310 mL/100 L water	
		NSW only	400 mL + 250 mL surfactant/ 100 L water		
		Qld only	75 mL/100 L water		
<b>Oil Tea Tree (<i>Melaleuca alternifolia</i>)</b>	Tip-Gall Midge ( <i>Dasineura</i> sp), Psyllids, Pyrgo Beetle	Qld, NSW only	340 mL/ha	5 months	Monitor the build up of Tip-Gall Midge in Spring by counting the trapped midge in spider webs. Spray when 10 percent of the growing points are showing the damaging effects of the Tip-Gall Midge larvae. <b>Boom Spray:</b> Apply in 50 - 100 L water/ha. <b>Aircraft:</b> Apply in 20 - 40 L water/ha. Rotate pyrethroid pesticides during Summer when spraying Pyrgo Beetle. Use methomyl products as the last seasonal spray for cleaning up any Apparent Decimator 400 Insecticide or pyrethroid resistant Pyrgo Beetles. Apply a maximum of 2 applications per crop growing cycle with a maximum of six weeks between applications.
<b>Duboisia</b>	Thrips	Qld, WA only	75 mL/100 L of water as an overall spray	-	Apply every 7 - 10 days or as pest population indicates.
<b>Wild Flowers, Proteas</b>	Aphids, Thrips Leafhoppers, Rutherglen Bug	WA only	75 mL/100 L of water	-	Apply when pests appear. Dimethoate will not control OP resistant mites.
<b>Trees: Eucalypts, Kurrajongs, Flame Trees, Umbrella Trees</b>	Jarrah Leaf Miner, Psyllids, Kurrajong Leaf Miner, Leaf Blister, Sawfly, Lerp Insects, Scale Insects, Spittle Bugs, Mites	WA only	25 mL/8 L of water	-	Apply in early June for control of Jarrah leaf miner and in early Spring for Psyllids. Trunk injection method may be used on large trees. Bore holes in tree trunk and fill with 1:1 mixtures of this product and water. Apply in late January as above for Kurrajong leaf miner.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**
**DO NOT USE THIS PRODUCT IN THE HOME GARDEN**
**WITHHOLDING PERIODS**

Litchi (pre-planting dip), Oilseeds (other than cotton and peanuts), Pasture forage crops: <b>NOT REQUIRED WHEN USED AS DIRECTED.</b>
Post Harvest Dipping (Avocados, Bananas, Cactus Fruit, Custard Apples, Feijoas, Guavas, Kiwifruit (Chinese Gooseberries inedible peel varieties), Lychees, Mangoes, Melons, Passionfruit, Banana, Passionfruit, Pawpaws, Pomegranates, Tamarillos): <b>NOT REQUIRED WHEN USED AS DIRECTED (DIP USES ONLY).</b>
Tomatoes, Large, Field Grown For Fresh Consumption: <b>NOT REQUIRED WHEN USED AS DIRECTED (IE. DO NOT APPLY AFTER COMMENCEMENT OF FLOWERING).</b>

**HARVEST WITHHOLDING PERIODS**

Blueberries (and other vaccinium berries including bilberries), Zucchini: <b>DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.</b>
Capsicums, Mango: <b>DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.</b>

Asparagus; Beans (green vegetables); Blackberries; Citrus; Melons (including watermelons); Onions; Peas (green vegetables, not snow or sugar snap peas); Raspberries; Rhubarb; Avocado, Litchi/Lychee: <b>DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.</b>
Beetroot, Cotton, Eggplant, Peanuts, Potatoes, Pulses (grain legumes), Sweet Potatoes, Turnip: <b>DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.</b>
Tomatoes (for processing): <b>DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.</b>
Cereals, (including maize, sorghum), Tobacco: <b>DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.</b>
Oil Tea Tree: <b>DO NOT HARVEST FOR 5 MONTHS AFTER APPLICATION.</b>
<b>GRAZING WITHHOLDING PERIODS</b>
Lucerne, Oilseeds (other than cotton and peanuts), Pastures, Pasture seed crops and Forage <b>NOT REQUIRED WHEN USED AS DIRECTED (i.e. DO NOT apply more than 7 days after crop emergence).</b>
Beans, Peas (green vegetables not snow or sugar snap peas): <b>DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 DAYS AFTER APPLICATION.</b>
Cereals, (Including Maize, Sorghum); Pulses (Grain Legumes): <b>DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION.</b>
Cotton: <b>DO NOT GRAZE OR CUT FOR STOCKFOOD.</b> <b>DO NOT FEED COTTON FODDER, STUBBLE OR TRASH TO LIVESTOCK.</b>

## GENERAL INSTRUCTIONS

### MIXING

The product can be poured directly into the water in the vat with agitators in operation. If combining with another product, mix each product separately in a small quantity of water first before adding to the vat.

### APPLICATION BY DILUTE SPRAYING

- Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

### APPLICATION BY CONCENTRATE SPRAYING

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

#### EXAMPLE ONLY

- Dilute spray volume as determined above: For example 1500 L/ha
  - Your chosen concentrate spray volume: For example 500 L/ha
  - The concentration factor in this example is:  $3 \times (\text{i.e. } 1500 \text{ L} \div 500 \text{ L} = 3)$
  - If the dilute label rate is 15 mL/100 L, then the concentrate rate becomes  $3 \times 15$  that is 45 mL/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

## REDLEGGED EARTH MITE

Redlegged Earth Mite (RLEM) is an introduced pasture and crop pest in southern Australia. RLEM is active in the cool wet months from May to November. During the 6 hotter months of the year RLEM avoid the hot dry conditions by developing a resting stage which is impervious to heat and drought. They do this by producing diapause (over-summering) eggs in Spring that remain on the soil surface. Very high numbers of over-summering eggs can be found on the soil surface, ready to emerge in the following Autumn, providing a threat to the germinating pasture or crop. The use of higher application rates in cereals and pasture after Autumn rains when mites emerge can provide extended periods of control. A system such as Timerite™ can also be used to estimate the optimum timing for a Spring spray to reduce egg-laying adult mite numbers and hence the damage to pasture and crops the following autumn when RLEM emerge from eggs.

### INSECTICIDE RESISTANCE WARNING

## GROUP 1B INSECTICIDE

For insect resistance management Apparent Decimator 400 Insecticide is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Apparent Decimator 400 Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Apparent Decimator 400 Insecticide or other Group 1B insecticides are used repeatedly. The effectiveness of Apparent Decimator 400 Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Decimator 400 Insecticide to control resistant insects. Apparent Decimator 400 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier or local agricultural department agronomist.

### RE-ENTRY AND RE-HANDLING

**Avocado, mango trees:** DO NOT allow entry into treated areas for 9 days for fruit thinning and for 2– days for hand harvesting. DO NOT allow entry into treated areas for hand pruning, irrigation, orchard maintenance, weeding, scouting, or transplanting until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

**Citrus trees:** DO NOT allow entry into treated areas for 4–days hand harvesting. DO NOT allow entry into treated areas for hand pruning, orchard maintenance, weeding, baiting/trapping, scouting, or transplanting until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

**Ornamentals-cut flowers or nursery plant:** DO NOT allow entry into treated areas for container moving, hand harvesting of cut flowers, hand irrigation, pinching, hand pruning, scouting, transplanting, and hand weeding until the spray has dried. If prior entry is required, wear

cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

**Ornamental trees farm and forest trees:** DO NOT allow entry into treated areas for 9 days for hand set irrigation. DO NOT allow entry into treated areas for 7–days for hand harvesting and for 1–day for hand pruning, shaping or scouting. DO NOT allow entry into treated areas for container moving, grading/tagging, transplanting or weeding until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

**Glasshouses and other confined areas:** DO NOT re-enter until spray deposits have dried and areas has been thoroughly ventilated.

**All other crops (litchi, blackberries, raspberries, vegetables, grain legumes, cereals, cotton, oilseeds, forage crops, tobacco, ornamental shrubs, duboisia, oil tea tree):** DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**Post-harvest dipping of fruit and vegetables, and pre-plant dipping of plants:** DO NOT handle treated fruit, vegetable or plant until the product solution has dried. If prior handling is required, wear elbow-length chemical resistant gloves.

### **PROTECTION OF LIVESTOCK**

Dangerous to bees. DO NOT spray any plants in flower while bees are actively foraging.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers, or waterways with the chemical or used containers. Dangerous to fish.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Protect from direct sunlight and temperatures above 40°C. If storing for periods of more than 2 - 3 months avoid temperatures above 30°C. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **DIP DISPOSAL**

Add 3 kg either slaked, hydrated or quick lime per 1000 litres of dip solution in a separate vessel to the dipping tank. Leave that mix for one or two hours to neutralise the chemical component. The inactivated mix can then be poured into a trench or sprayed on grass. DO NOT flush to rivers, creeks or drain ways.

### **SAFETY DIRECTIONS**

Poisonous if absorbed by skin contact or inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Will damage eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, a PVC or rubber apron, elbow-length chemical resistant gloves, face shield and impervious footwear. When using the prepared spray (or dip for pre-plant and post-harvest dipping) wear elbow-length chemical resistant gloves. If applying by hand by vehicle mounted low pressure equipment wear cotton overalls buttoned to the neck and wrist, elbow-length chemical resistant gloves and a half face-piece respirator with organic vapour/gas cartridge or canister. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face-shield, respirator and contaminated clothing.

### **FIRST AID**

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre. Phone Australia 13 11 26 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet available from the supplier.

### **CONDITIONS OF SALE**

The use of Apparent Decimator 400 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.