# **POISON**

**KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING** 



# Pilferer 500

# **HERBICIDE**

**ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE** SOLVENT: 42 g/L ETHYLENE GLYCOL



HERBICIDE

For selective control of certain grasses and broad leaf weeds in Lettuce, Sports Turf, Home Lawns and Legume Seed Crops and Pastures, as per the Direction for Use.











**DIRECTIONS FOR USE** 

Situation	Weeds Controlled	State	Rate L/ha	Critical Comments		
Lettuce	Barnyard Grass, Blackberry Nightshade,	All	4.5 in	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.		
(direct sown)	Chickweed, Portulaca, Winter Grass, Ryegrass,	States	330 – 550 L			
Transplanted	Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	/ initial activities and activities and initial activities and activities and activities and activities activities activities and activities activities activities activities and activities		ely after transplanting. For best results, prepare the seed bed free from clods. ion to wet the top 3 to 4 cm of soil should be applied within 24 hours.		
	0		2.3 in	Apply on light sandy soils, low in organic matter.		
			300 – 550 L			
Turf & Lawns	Winter grass	All	1.2	Rate/ Bowling	Apply during the autumn and Winter, if established turf is infested with	
Common	3	States	(12 mL /100m <sup>2</sup> )		seedling winter grass. Ensure a uniform coverage. Irrigate immediately	
Couch			,	37 x 37m	after application with 12 - 25 mm irrigation or rain and again 24 hours	
				170 mL	later.	
Oueensland		Old	0.6	85 mL	Repeat if necessary on Qld. Blue Couch after 4 weeks.	
blue couch		only	(6 mL /100m <sup>2</sup> )	05 1112	DO NOT re-seed for 60 days after treatment.	
Buffalo grass		WA	1.2	170 mL	Use ONLY on established turf.	
Danaio grass		only	(12 mL /100m <sup>2</sup> )		<b>DO NOT</b> use on Bluegrass, ryegrass, fescue or bent grass lawns.	
Lucerne,	Annual Ryegrass, Winter grass, Barley grass,	NSW,	1 - 1.5	SEEDLING CR	<b>EEDLING CROPS:</b> Apply after the third trifoliate leaf is fully developed.	
Clover,	Silver Grass, Prairie Grass, Canary Grass, Great	ACT,		ESTABLISHED CROPS: Apply any time before the end of July.		
Medics etc.	Brome, Squirrel Tail Fescue and Rats Tail Fescue   SA		SA,	WEEDS: Apply when weeds are at post or pre-emergent stages.  BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the		
Grown for	Wild Oats, Perennial Ryegrass, Bentgrass, English	grass, English WA, 1.5 - 2				
forage, Hay	Couch, Yorkshire Fog Grass, Chickweed, seedling		_			
or Seed	Sorrel, Blackberry Nightshade, Patterson's Curse,	only		higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatments is recommended.		
production	seedling Wireweed	-				

## NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

#### WITHHOLDING PERIOD:

#### DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

#### GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet

Prior to pouring, shake container vigorously, then add the required quantity of Apparent Pilferer 500 Herbicide to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

### RESISTANT WEEDS WARNING GROUP | HERBICIDE

Apparent Pilferer 500 Herbicide is a member of the benzamide group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management Apparent Pilferer 500 Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to Apparent Pilferer 500 Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Pilferer 500 Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from failure of Apparent Pilferer 500 Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be used can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Apparent Pty Ltd representative.
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND

## ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

APVMA Approval No: 68267/102232 Batch No:

#### PROTECTION OF LIVESTOCK

DO NOT graze sheep on treated areas or feed to poultry or

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, wellventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and win inflate the eyes and skin. Avoid Contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Date of Manufacture:

#### SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier. CONDITIONS OF SALE

The use of Apparent Pilferer 500 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture: Suspected of causing cancer. Very toxic to aquatic life. Very toxic to aquatic life with long lasting effects. Avoid release to the environment. IF ON SKIN: Wash with plenty of soap and water. IF exposed or concerned: Get medical advice/ attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing If skin irritation occurs: Get medical advice/ attention. Take off contaminated clothing and wash before reuse. Collect spillage. Store locked up

> Scan QR code to view booklet on mobile device