

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent



Flutriafol 25/Cypermethrin 4

LIQUID SEED DRESSING

ACTIVE CONSTITUENTS: 25 g/L FLUTRIAFOL
4 g/L CYPERMETHRIN

SOLVENT: 488 g/L ETHYLENE GLYCOL

GROUP 3 FUNGICIDE

GROUP 3A INSECTICIDE

Seed dressing treatment to protect against smuts and bunt in wheat, barley, oats and triticale.

PROUDLY



OWNED



The
Choice is
Apparent



CONTENTS

10

LITRES

APPARENT PTY LTD A.C.N. 143 724 136

Suite G.08 | 762 Toorak Road, Hawthorn East VIC 3123 Australia

T 03 9822 1321 M 0411 227 338 E enquiries@apparentag.com.au www.apparentag.com.au

DIRECTIONS FOR USE

RESTRAINT: DO NOT apply "Vydate" to seed pre-treated with this product.

CROP	DISEASE CONTROLLED	STATE	RATE/100 kg SEED	CRITICAL COMMENTS
Wheat	Bunt (<i>Tilletia foetida</i> , <i>T. caries</i>), Loose Smut (<i>Ustilago segetum</i> var <i>tritici</i>), Soilborne Flag Smut (<i>Urocystis agropyri</i>), Seedeborne Flag Smut (<i>Urocystis agropyri</i>)	All States	100 mL	This product is to be applied to grain undiluted. Seed must be treated to ensure even coverage of all grains, using correctly calibrated seed treatment equipment. Reduced emergence may occur if seed treated with Flutriafol is sown to a depth greater than 5 cm or into soil at a temperature less than 5°C or greater than 20°C.
Barley	Covered Smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose Smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Oats	Covered Smut (<i>Ustilago segetum</i> var <i>segetum</i>), Loose Smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Triticale	Loose Smut (<i>Ustilago segetum</i> var <i>tritici</i>)			
Wheat, Barley, Oats & Triticale	Protection against insect pests of stored grains: Lesser Grain Borer (<i>Phytophthora dominica</i>), Granary Weevil (<i>Stophilus granarius</i>), Rice Weevil (<i>Stophilus oryzae</i>), Rust-red Flour Beetle (<i>Tribolium castaneum</i>), Saw Toothed Grain Beetle (<i>Oryzaephilus surinamensis</i>), Tropical Warehouse Moth (<i>Ephestia cautella</i>)			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 4 WEEKS OF SOWING.

GENERAL INSTRUCTIONS

Seed may be treated with Apparent Flutriafol 25 / Cypermethrin 4 Liquid Seed Dressing at any time from harvest to sowing. Treated seed may be held over for sowing in the following season. The product contains Cypermethrin for protection against insect pests of stored grain.

THIS PRODUCT IS NOT MISCIBLE WITH WATER AND CANNOT BE DILUTED.

Equipment must be dry.

CLEANING UP

Equipment should be thoroughly cleaned with methylated spirits, not water.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Apparent Flutriafol 25/Cypermethrin 4 Liquid Seed Dressing is a member of the DMJ group of fungicides. For fungicide resistance management, the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

INSECTICIDE RESISTANCE WARNING

GROUP 3A INSECTICIDE

For insecticide resistance management Apparent Flutriafol 25/Cypermethrin 4 Liquid

Seed Dressing is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to the product and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if this product or other Group 3A insecticides are used repeatedly. The effectiveness of this product on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects.

Apparent Flutriafol 25/Cypermethrin 4 Liquid Seed Dressing may be subject to specific resistance management strategies. For further information contact your local supplier, Apparent Pty Ltd representative or local agricultural department agronomist.

PROTECTION OF CROPS, NATURE AND OTHER NON-TARGET PLANTS

Short coleoptile cuttings should not be sown too deep. In some situations, trifluralin herbicides and triazole fungicides interact to suppress emergence, if possible avoid this contamination. Any seed not intended to be used for sowing should be destroyed.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or be allowed to contaminate seed intended for animal consumption. Seed should not be left exposed on the ground (particularly during sowing). Destroy treated seed not intended for sowing.

PROTECTION OF WILDLIFE, FISH, CRUSTACEAN AND THE ENVIRONMENT

DO NOT feed or expose treated seed to wildlife and domestic animals, particularly birds.

Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical, or containers which have held the chemical or treated seed.

STORAGE AND DISPOSAL

Seed treated with this product must not be used for human consumption or be allowed to contaminate seed intended for human consumption. Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with this product. Containers which have held treated seed should not be used for any other purpose.

For Non-Refillable Containers: Store product in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Product is poisonous if swallowed. May

irritate the eyes and skin. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. DO NOT inhale dust, vapour or spray mist. When preparing and using the product wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE: The use of Apparent Flutriafol 25/Cypermethrin 4 Liquid Seed Dressing beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

