

blank page

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent 

Swoosh 125 SC

FUNGICIDE

ACTIVE CONSTITUENT: 125g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf rust, stripe rust, and septoria nodorum blotch of wheat and leaf rust of barley, as specified in the Directions for Use Table.

IMPORTANT: Read the attached booklet before use.

APVMA Approval No: 67738/102432

AIRR APPARENT PTY LTD
15/16 Princes Street, Newport NSW 2106
M 0411 227 338 | E enquiries@apparentag.com.au | www.apparentag.com.au

DIRECTIONS FOR USE

CROP	DISEASE	RATE PER HA
Wheat	Leaf rust (<i>Puccinia recondite</i>)	500 mL plus
	Stripe rust (<i>Puccinia striiformis</i>)	200 mL Chemwet per 100 L of spray mix
	Septoria nodorum blotch (<i>Septoria nodorum</i>)	500 mL plus 200 mL Chemwet per 100 L of spray mix
Barley	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL plus 200 mL Chemwet per 100 L of spray mix

CRITICAL COMMENTS

Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32 - 59). DO NOT apply later than ZGS 59.

Repeat spraying may be required for stripe rust. Regularly monitor crops from 3 – 4 weeks after spraying for signs of re-infection and treat accordingly.

Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32 – 59). DO NOT apply later than ZGS 59.

Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Apparent Swoosh 125 SC Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to the new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS 39) may not provide any yield benefit, particularly under dry seasonal conditions.

MIXING

Apparent Swoosh 125 SC Fungicide is a suspension concentrate formulation. Add the product to the half filled spray vat while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground: Apply in a water volume of between 50 and 100 L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial: Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20 L/ha.

COMPATIBILITY

For information on compatibility please contact your local reseller or AIRR Apparent Pty Ltd representative.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Apparent Swoosh 125 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Apparent Swoosh 125 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried.

CAUTION

Export of treated cereals: Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with Apparent Swoosh 125 SC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with AIRR Apparent Pty Ltd or your grain exporter for the latest information on MRLs and import tolerances BEFORE using Apparent Swoosh 125 SC Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI)

AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, crop land, livestock, pastures, wetlands, waterbodies or watercourses. Do NOT apply Apparent Swoosh 125 SC Fungicide aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and halpiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which is available from the supplier.

CONDITIONS OF SALE

The use of Apparent Swoosh 125 SC Fungicide being beyond the control of the manufacturer no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AIRR Apparent Pty Ltd accepts with no responsibility for any consequence whatsoever resulting from the use of this product.